arXiv:math/0105024v3 [math.RT] 15 May 2001

A Twining Character Formula for Demazure Modules

By

Daisuke Sagaki
Graduate School of Mathematics, University of Tsukuba,
Tsukuba, Ibaraki 305-8571, Japan (e-mail: sagaki@math.tsukuba.ac.jp)

In [FRS] and [FSS], they introduced new character-like quantities corresponding to a graph automorphism of a Dynkin diagram, called twining characters, for certain Verma modules and integrable highest weight modules over a symmetrizable Kac-Moody algebra, and gave twining character formulas for them. Recently, the notion of twining characters has naturally been extended to various modules, and formulas for them has been given ([KN], [KK], [N1]–[N4]). In this short note, we introduce a twining character formula for Demazure modules over a symmetrizable Kac-Moody algebra, which is obtained in [S].

Let $\mathfrak{g} = \mathfrak{g}(A) = \mathfrak{n}_- \oplus \mathfrak{h} \oplus \mathfrak{n}_+$ be a symmetrizable Kac-Moody algebra over $\mathbb{Q}$ associated to a GCM $A = (a_{ij})_{i,j \in I}$ of finite size, where $\mathfrak{h}$ is the Cartan subalgebra, $\mathfrak{n}_+$ the sum of positive root spaces, and $\mathfrak{n}_-$ the sum of negative root spaces, and let $\omega : I \to I$ be a (Dynkin) diagram automorphism, that is, a bijection $\omega : I \to I$ satisfying $a_{\omega(i),\omega(j)} = a_{ij}$ for all $i, j \in I$. It is known that a diagram automorphism induces a Lie algebra automorphism $\omega \in \mathrm{Aut}(\mathfrak{g})$ that preserves the triangular decomposition of $\mathfrak{g}$ (see [FSS] and [FRS]). Then we define a linear automorphism $\omega^* \in \mathrm{GL}(\mathfrak{h}^*)$ by $(\omega^*(\lambda))(h) := \lambda(\omega(h))$ for $\lambda \in \mathfrak{h}^*$, $h \in \mathfrak{h}$. We set $(\mathfrak{h}^*)^0 := \{\lambda \in \mathfrak{h}^* \,|\, \omega^*(\lambda) = \lambda\}$, and call its elements symmetric weights. We also set $\widetilde{W} := \{w \in W \,|\, w\,\omega^* = \omega^*\,w\}$.

Further we define a "folded" matrix $\widehat{A}$ associated to ω, which is again a symmetrizable GCM if ω satisfies a certain condition, called the linking condition (see [FSS]). Then the Kac-Moody algebra $\widehat{\mathfrak{g}} = \mathfrak{g}(\widehat{A})$ associated to $\widehat{A}$ is called the orbit Lie algebra. We denote by $\widehat{\mathfrak{h}}$ the Cartan subalgebra of $\widehat{\mathfrak{g}}$ and by $\widehat{W}$ the Weyl group of $\widehat{\mathfrak{g}}$. Then there exist a linear isomorphism $P_\omega^* : \widehat{\mathfrak{h}}^* \to (\mathfrak{h}^*)^0$ and a group isomorphism $\Theta : \widehat{W} \to \widetilde{W}$ such that $\Theta(\widehat{w}) = P_\omega^* \circ \widehat{w} \circ (P_\omega^*)^{-1}$ for all $\widehat{w} \in \widehat{W}$.

Let λ be a dominant integral weight. Denote by $L(\lambda) = \bigoplus_{\chi \in \mathfrak{h}^*} L(\lambda)_\chi$ the irreducible highest weight $\mathfrak{g}$-module of highest weight λ. Then, for $w \in W$, we define the Demazure module $L_w(\lambda)$ of lowest weight $w(\lambda)$ in $L(\lambda)$ by $L_w(\lambda) := U(\mathfrak{b})u_{w(\lambda)}$, where $u_{w(\lambda)} \in L(\lambda)_{w(\lambda)} \setminus \{0\}$ and $U(\mathfrak{b})$ is the universal enveloping algebra of the Borel subalgebra $\mathfrak{b} :=$

$\mathfrak{h} \oplus \mathfrak{n}_+$ of $\mathfrak{g}$. If λ is symmetric, then we know from [FSS] and [FRS] (see also [N1]) that there exists a (unique) linear automorphism $\tau_\omega : L(\lambda) \to L(\lambda)$ such that

$$\tau_\omega(xv) = \omega^{-1}(x)\tau_\omega(v) \quad \text{for all } x \in \mathfrak{g},\ v \in L(\lambda)$$

and $\tau_\omega(u_\lambda) = u_\lambda$ with u_λ a (nonzero) highest weight vector of $L(\lambda)$. Then it is easily seen that the Demazure module $L_w(\lambda)$ with $w \in \widetilde{W}$ is τ_ω-stable. Here we define the twining character $\mathrm{ch}^\omega(L_w(\lambda))$ of $L_w(\lambda)$ by:

$$\mathrm{ch}^\omega(L_w(\lambda)) := \sum_{\chi \in (\mathfrak{h}^*)^0} \mathrm{tr}\big(\tau_\omega|_{L_w(\lambda)_\chi}\big) e(\chi).$$

Our formula is the following:

Theorem. *Let* λ *be a symmetric dominant integral weight and* $w \in \widetilde{W}$. *Set* $\widehat{\lambda} := (P_\omega^*)^{-1}(\lambda)$ *and* $\widehat{w} := \Theta^{-1}(w)$. *Then we have*

$$\mathrm{ch}^\omega(L_w(\lambda)) = P_\omega^*(\mathrm{ch}\,\widehat{L}_{\widehat{w}}(\widehat{\lambda})),$$

where $\widehat{L}_{\widehat{w}}(\widehat{\lambda})$ *is the Demazure module of lowest weight* $\widehat{w}(\widehat{\lambda})$ *in the irreducible highest weight module* $\widehat{L}(\widehat{\lambda})$ *of highest weight* $\widehat{\lambda}$ *over the orbit Lie algebra* $\widehat{\mathfrak{g}}$.

REFERENCES

[FRS] J. Fuchs, U. Ray, and C. Schweigert, *Some automorphisms of generalized Kac-Moody algebras*, J. Algebra **191** (1997), 518–540.

[FSS] J. Fuchs, B. Schellekens, and C. Schweigert, *From Dynkin diagram symmetries to fixed point structures*, Comm. Math. Phys. **180** (1996), 39–97.

[KN] M. Kaneda and S. Naito, *A twining character formula for Demazure modules*, preprint.

[KK] S.-J. Kang and J.-H. Kwon, *Graded Lie superalgebras, supertrace formula, and orbit Lie superalgebras*, Proc. London Math. Soc. **81** (2000), 675–724.

[N1] S. Naito, *Twining character formula of Kac-Wakimoto type for affine Lie algebras*, preprint.

[N2] ———, *Twining characters and Kostant's homology formula*, preprint.

[N3] ———, *Twining characters, Kostant's homology formula, and the Bernstein-Gelfand-Gelfand resolution*, preprint.

[N4] ———, *Twining character formula of Borel-Weil-Bott type*, preprint.

[S] ———, *Crystal bases, path models, and a twining character formula for Demazure modules*, preprint.